

The F- and t-Distributions

Suppose U_1, U_2 are two independent random variables and

$$U_1 \sim \chi_{p_1}^2, U_2 \sim \chi_{p_2}^2.$$

Then

$$F = \frac{U_1/p_1}{U_2/p_2}$$

has the F-Distribution with p_1 and p_2 DF, denoted by

$$F \sim F_{p_1, p_2}.$$

Result 5.12:

The density of $F \sim F_{p_1, p_2}$ is

$$f_F(t) = \frac{\Gamma\left(\frac{p_1+p_2}{2}\right) \left(\frac{p_1}{p_2}\right)^{\frac{p_1}{2}}}{\Gamma\left(\frac{p_1}{2}\right) \Gamma\left(\frac{p_2}{2}\right)} t^{\frac{p_1}{2}-1} \left(1 + \frac{p_1}{p_2}t\right)^{-\frac{p_1+p_2}{2}}.$$

Proof of Result 5.12:

HW problem. □

Suppose U_1 and U_2 are independent random variables and suppose

$$U_1 \sim \chi_{p_1}^2(\phi) \quad \text{and} \quad U_2 \sim \chi_{p_2}^2.$$

Then

$$F = \frac{U_1/p_1}{U_2/p_2}$$

has the noncentral F -distribution with p_1 and p_2 DF and NCP ϕ .

$$(F \sim F_{p_1, p_2}(\phi))$$

Result 5.13:

Suppose $W \sim F_{p_1, p_2}(\phi)$. Then for fixed p_1, p_2 and $c > 0$, $\mathbb{P}(W > c)$ is a strictly increasing function of ϕ .

Proof:

$$W \stackrel{d}{=} \frac{U_1/p_1}{U_2/p_2},$$

where U_1 independent of U_2 , $U_1 \sim \chi_{p_1}^2(\phi)$, and $U_2 \sim \chi_{p_2}^2$.

Thus,

$$\begin{aligned}\mathbb{P}(W > c) &= \mathbb{P}_\phi(W > c) \\ &= \mathbb{P}_\phi\left(\frac{U_1/p_1}{U_2/p_2} > c\right) \\ &= \mathbb{P}_\phi\left(U_1 > \frac{cp_1}{p_2}U_2\right) \\ &= \int_0^\infty \mathbb{P}_\phi\left(U_1 > \frac{cp_1}{p_2}u_2 \mid U_2 = u_2\right) f_{U_2}(u_2) du_2\end{aligned}$$

$$= \int_0^{\infty} g_{\phi}(u_2) f_{U_2}(u_2) du_2$$

where

$$\begin{aligned} g_{\phi}(u_2) &= \mathbb{P}_{\phi} \left(U_1 > \frac{c p_1}{p_2} u_2 \mid U_2 = u_2 \right) \\ &= \mathbb{P}_{\phi} \left(U_1 > \frac{c p_1}{p_2} u_2 \right) \quad \text{by ind. of } U_1, U_2. \end{aligned}$$

By Result 5.11,

$$g_{\phi_1}(u_2) < g_{\phi_2}(u_2) \quad \forall 0 \leq \phi_1 < \phi_2 \quad \text{and} \quad \forall u_2 > 0.$$

Thus,

$$\begin{aligned} 0 &< \int_0^{\infty} (g_{\phi_2}(u_2) - g_{\phi_1}(u_2))f_{U_2}(u_2)du_2 \\ &= \int_0^{\infty} g_{\phi_2}(u_2)f_{U_2}(u_2)du_2 - \int_0^{\infty} g_{\phi_1}(u_2)f_{U_2}(u_2)du_2 \\ &= \mathbb{P}_{\phi_2}(W > c) - \mathbb{P}_{\phi_1}(W > c) \quad \forall 0 \leq \phi_1 < \phi_2. \end{aligned}$$

$\therefore \mathbb{P}_{\phi}(W > c)$ is a strictly increasing function of ϕ . □

Suppose

$$U \sim N(\mu, 1) \quad \text{and} \quad V \sim \chi_k^2.$$

If U and V are independent, then

$$T = \frac{U}{\sqrt{V/k}}$$

has the noncentral t -distribution with k DF and NCP μ . ($T \sim t_k(\mu)$)

If $\mu = 0$, then $T = U/\sqrt{V/k}$ has Student's t -distribution with k degree of freedom and density

$$f_T(t) = \frac{\Gamma\left(\frac{k+1}{2}\right)}{\Gamma\left(\frac{k}{2}\right)\sqrt{\pi k}} \left(1 + \frac{t^2}{k}\right)^{-\frac{k+1}{2}}.$$

We use $T \sim t_k$ to indicate that T has Student's t -distribution with k DF.

Suppose $T \sim t_k(\mu)$.

Find the distribution of T^2 .